

International customers can order Q-potencies in powder form. Each packet of powder contains one globule of the desired Q-potency in a lactose base. This globule can then be diluted by the doctor in different ways. We recommend using a 10 ml bottle with a water/alcohol (at least 20%) mixture. Make sure that room is left in the bottle for shaking required to potentize the solution.

Instructions for the Administration of Q-potencies


Assuming that the physician has already advised the patient of the number drops to be used and at what intervals, the physician (or patient) should then prepare the remedy in following manner:

1. The first dose:



Place the number of drops (we recommend 2) prescribed by the physician into a plastic tumbler (capacity of 0,2 – 0,3l) filled to 2/3 with cold water and stir 10 times with a plastic spoon. Take one teaspoon of this solution and discard the remainder. Quickly rinse the spoon and tumbler with water. Do not wash with other dishes.

2. Second and all subsequent doses:



Shake the bottle containing the remedy vigorously 10 times, place the number of drops prescribed by your physician into the previously used tumbler filled, as above, with water, stir 10 times and take one teaspoon of it, discarding the rest. Rinse tumbler and spoon as before.

At the end of the prescribed time for this remedy, discard the spoon, the tumbler and the remainder of the remedy. The same procedure should be followed whenever your physician prescribes a remedy.

Homöopathisches Labor Gudjons
Höfatsweg 21, D-86391 Stadtbergen-Deuringen
Telefon +49 821 4447877 - Fax +49 821 438444
e-mail: labor@gudjons.com - www.gudjons.com